

All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books

Search for

Display Show Sort by Send to

About Entrez
NCBI Toolbar

All: 6 Review: 0

Items 1 - 6 of 6

One page.

Text Version

☐ 1: [Okuda D, Horii K, Mizuno H, Morita T.](#) Related Articles, Links

Characterization and preliminary crystallographic studies of EMS16, an antagonist of collagen receptor (GPIa/IIa) from the venom of *Echis multisquamatus*.
J Biochem (Tokyo). 2003 Jul;134(1):19-23.
PMID: 12944366 [PubMed - indexed for MEDLINE]

Entrez PubMed
Overview
Help | FAQ
Tutorials
New/Noteworthy
E-Utilities

☐ 2: [Augustin A, Muller-Steffner H, Schuber F.](#) Related Articles, Links

Molecular cloning and functional expression of bovine spleen ecto-NAD⁺ glycohydrolase: structural identity with human CD38.
Biochem J. 2000 Jan 1;345 Pt 1:43-52.
PMID: 10600637 [PubMed - indexed for MEDLINE]

PubMed Services
Journals Database
MeSH Database
Single Citation
Matcher
Batch Citation
Matcher
Clinical Queries
Special Queries
LinkOut
My NCBI

☐ 3: [Bourrat C, Radisson J, Chavassieux P, Azzar G, Roux B, Meunier PJ.](#) Related Articles, Links

Activity increase after extraction of alkaline phosphatase from human osteoblastic membranes by nonionic detergents: influence of age and sex.
Calcif Tissue Int. 2000 Jan;66(1):22-8.
PMID: 10602840 [PubMed - indexed for MEDLINE]

☐ 4: [Del Vecchio Blanco F, Bolognesi A, Malorni A, Sande MJ, Savino G, Parente A.](#) Related Articles, Links

Complete amino-acid sequence of PD-S2, a new ribosome-inactivating protein from seeds of *Phytolacca dioica* L.
Biochim Biophys Acta. 1997 Mar 7;1338(1):137-44.
PMID: 9074624 [PubMed - indexed for MEDLINE]

☐ 5: [Munishkin A, Wool IG.](#) Related Articles, Links

Systematic deletion analysis of ricin A-chain function. Single amino acid deletions.
J Biol Chem. 1995 Dec 22;270(51):30581-7.
PMID: 8530493 [PubMed - indexed for MEDLINE]

Related
Resources
Order Documents
NLM Mobile
NLM Catalog
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

6: [Yoshino M, Takagi T.](#)

[Related Articles, Links](#)



Subunit structure of AMP nucleosidase from *Azotobacter vinelandii*.
J Biochem (Tokyo). 1973 Dec;74(6):1151-6. No abstract available.
PMID: 4781054 [PubMed - indexed for MEDLINE]

Display [Abstract](#) Show [20](#) Sort by [Send to](#)


[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Mar 28 2006 06:29:55



All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books

Search for

Limits Preview/Index History Clipboard Details

Display Show Sort by Send to

About Entrez
NCBI Toolbar

All: 13 Review: 0 ☐

Items 1 - 13 of 13

One page.

Text Version

☐ 1: [Nakamura I, Yoneda H, Maeda T, Makino A, Ohmae M, Sugiyama J, Ueda M, Kobayashi S, Kimura S.](#) Related Articles, Links

☐ Enzymatic polymerization behavior using cellulose-binding domain deficient endoglucanase II.
Macromol Biosci. 2005 Jul 14;5(7):623-8.
PMID: 15988789 [PubMed - indexed for MEDLINE]

☐ 2: [Zhou W, Irwin DC, Escovar-Kousen J, Wilson DB.](#) Related Articles, Links

☐ Kinetic studies of Thermobifida fusca Cel9A active site mutant enzymes.
Biochemistry. 2004 Aug 3;43(30):9655-63.
PMID: 15274620 [PubMed - indexed for MEDLINE]

☐ 3: [Mandelman D, Belaich A, Belaich JP, Aghajari N, Driguez H, Haser R.](#) Related Articles, Links

☐ X-Ray crystal structure of the multidomain endoglucanase Cel9G from Clostridium cellulolyticum complexed with natural and synthetic cello-oligosaccharides.
J Bacteriol. 2003 Jul;185(14):4127-35.
PMID: 12837787 [PubMed - indexed for MEDLINE]

☐ 4: [Kataeva IA, Seidel RD 3rd, Shah A, West LT, Li XL, Ljungdahl LG.](#) Related Articles, Links

☐ The fibronectin type 3-like repeat from the Clostridium thermocellum cellobiohydrolase CbhA promotes hydrolysis of cellulose by modifying its surface.
Appl Environ Microbiol. 2002 Sep;68(9):4292-300.
PMID: 12200278 [PubMed - indexed for MEDLINE]

☐ 5: [Saloheimo M, Paloheimo M, Hakola S, Pere J, Swanson B, Nyyssonen E, Bhatia A, Ward](#) Related Articles, Links

Related
Resources
Order Documents
NLM Mobile
NLM Catalog
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

M. Penttila M.



Swollenin, a *Trichoderma reesei* protein with sequence similarity to the plant expansins, exhibits disruption activity on cellulosic materials.

Eur J Biochem. 2002 Sep;269(17):4202-11.

PMID: 12199698 [PubMed - indexed for MEDLINE]

☐ 6: Ding SY, Bayer EA, Steiner D, Shoham Y, Lamed R. Related Articles, Links



A scaffoldin of the *Bacteroides cellulosolvens* cellulosome that contains 11 type II cohesins.

J Bacteriol. 2000 Sep;182(17):4915-25.

PMID: 10940036 [PubMed - indexed for MEDLINE]

☐ 7: Riedel K, Ritter J, Bauer S, Bronnenmeier K. Related Articles, Links



The modular cellulase CelZ of the thermophilic bacterium *Clostridium stercorarium* contains a thermostabilizing domain.

FEMS Microbiol Lett. 1998 Jul 15;164(2):261-7.

PMID: 9682475 [PubMed - indexed for MEDLINE]

☐ 8: Zverlov V, Mahr S, Riedel K, Bronnenmeier K. Related Articles, Links



Properties and gene structure of a bifunctional cellulolytic enzyme (CelA) from the extreme thermophile '*Anaerocellum thermophilum*' with separate glycosyl hydrolase family 9 and 48 catalytic domains.

Microbiology. 1998 Feb;144 (Pt 2):457-65.

PMID: 9493383 [PubMed - indexed for MEDLINE]

☐ 9: Sakon J, Irwin D, Wilson DB, Karplus PA. Related Articles, Links



Structure and mechanism of endo/exocellulase E4 from *Thermomonospora fusca*.

Nat Struct Biol. 1997 Oct;4(10):810-8.

PMID: 9334746 [PubMed - indexed for MEDLINE]

☐ 10: Pages S, Gal L, Belaich A, Gaudin C, Tardif C, Belaich JP. Related Articles, Links



Role of scaffolding protein CipC of *Clostridium cellulolyticum* in cellulose degradation.

J Bacteriol. 1997 May;179(9):2810-6.

PMID: 9139893 [PubMed - indexed for MEDLINE]

☐ 11: Macarron R, Henrissat B, Claeysens M. Related Articles, Links



Family A cellulases: two essential tryptophan residues in endoglucanase III from *Trichoderma reesei*.

Biochim Biophys Acta. 1995 Oct 19;1245(2):187-90.

PMID: 7492576 [PubMed - indexed for MEDLINE]

- ☐ 12: [Lassig JP, Shultz MD, Gooch MG, Evans BR, Woodward J.](#) Related Articles, Links



Inhibition of cellobiohydrolase I from *Trichoderma reesei* by palladium.

Arch Biochem Biophys. 1995 Sep 10;322(1):119-26.

PMID: 7574665 [PubMed - indexed for MEDLINE]

- ☐ 13: [Goldstein MA, Doi RH.](#) Related Articles, Links



Mutation analysis of the cellulose-binding domain of the *Clostridium cellulovorans* cellulose-binding protein A.

J Bacteriol. 1994 Dec;176(23):7328-34.

PMID: 7961505 [PubMed - indexed for MEDLINE]

Display Show Sort by Send to

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Mar 28 2006 06:29:55

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSSPTA1623SQS

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *
* *

NEWS 1		Web Page URLs for STN Seminar Schedule - N. America
NEWS 2		"Ask CAS" for self-help around the clock
NEWS 3	DEC 21	IPC search and display fields enhanced in CA/Caplus
with the		
		IPC reform
NEWS 4	DEC 23	New IPC8 SEARCH, DISPLAY, and SELECT fields in
USPATFULL/		
		USPAT2
NEWS 5	JAN 13	IPC 8 searching in IFIPAT, IFIUDB, and IFICDB
NEWS 6	JAN 13	New IPC 8 SEARCH, DISPLAY, and SELECT enhancements
added to		
		INPADOC
NEWS 7	JAN 17	Pre-1988 INPI data added to MARPAT
NEWS 8	JAN 17	IPC 8 in the WPI family of databases including WPIFV
NEWS 9	JAN 30	Saved answer limit increased
NEWS 10	JAN 31	Monthly current-awareness alert (SDI) frequency
added to TULSA		
NEWS 11	FEB 21	STN AnaVist, Version 1.1, lets you share your STN
AnaVist		
		visualization results
NEWS 12	FEB 22	Status of current WO (PCT) information on STN
NEWS 13	FEB 22	The IPC thesaurus added to additional patent
databases on STN		
NEWS 14	FEB 22	Updates in EPFULL; IPC 8 enhancements added
NEWS 15	FEB 27	New STN AnaVist pricing effective March 1, 2006
NEWS 16	FEB 28	MEDLINE/LMEDLINE reload improves functionality
NEWS 17	FEB 28	TOXCENTER reloaded with enhancements
NEWS 18	FEB 28	REGISTRY/ZREGISTRY enhanced with more experimental
spectral		
		property data
NEWS 19	MAR 01	INSPEC reloaded and enhanced
NEWS 20	MAR 03	Updates in PATDPA; addition of IPC 8 data without
attributes		
NEWS 21	MAR 08	X.25 communication option no longer available after
June 2006		

NEWS 22 MAR 22 EMBASE is now updated on a daily basis
NEWS 23 APR 03 New IPC 8 fields and IPC thesaurus added to
PATDPAFULL
NEWS 24 APR 03 Bibliographic data updates resume; new IPC 8 fields
and IPC
thesaurus added in PCTFULL

NEWS EXPRESS FEBRUARY 15 CURRENT VERSION FOR WINDOWS IS V8.01a,
CURRENT MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP),
AND CURRENT DISCOVER FILE IS DATED 19 DECEMBER 2005.
V8.0 AND V8.01 USERS CAN OBTAIN THE UPGRADE TO V8.01a

AT

<http://download.cas.org/express/v8.0-Discover/>

NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS LOGIN Welcome Banner and News Items

Enter NEWS followed by the item number or name to see news on that
specific topic.

All use of STN is subject to the provisions of the STN Customer
agreement. Please note that this agreement limits use to scientific
research. Use for software development or design or implementation
of commercial gateways or other similar uses is prohibited and may
result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *
* *

FILE 'HOME' ENTERED AT 12:26:33 ON 03 APR 2006

=> File Medline EMBASE Biosis Caplus AGRICOLA CABA

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.21	0.21

FILE 'MEDLINE' ENTERED AT 12:26:58 ON 03 APR 2006

FILE 'EMBASE' ENTERED AT 12:26:58 ON 03 APR 2006
Copyright (c) 2006 Elsevier B.V. All rights reserved.

FILE 'BIOSIS' ENTERED AT 12:26:58 ON 03 APR 2006
Copyright (c) 2006 The Thomson Corporation

FILE 'CAPLUS' ENTERED AT 12:26:58 ON 03 APR 2006
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2006 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'AGRICOLA' ENTERED AT 12:26:58 ON 03 APR 2006

FILE 'CABA' ENTERED AT 12:26:58 ON 03 APR 2006

COPYRIGHT (C) 2006 CAB INTERNATIONAL (CABI)

=> s (carbohydrate binding domain) or (CBD)
L1 8600 (CARBOHYDRATE BINDING DOMAIN) OR (CBD)

=> S L1 (6a) (structure)
L2 178 L1 (6A) (STRUCTURE)

=> s l1 (w) III
L3 0 L1 (W) III

=> l2 and review
L2 IS NOT A RECOGNIZED COMMAND
The previous command name entered was not recognized by the system.
For a list of commands available to you in the current file, enter
"HELP COMMANDS" at an arrow prompt (=>).

=> s l2 and review
L4 9 L2 AND REVIEW

=> duplicate
ENTER REMOVE, IDENTIFY, ONLY, OR (?):remove
ENTER L# LIST OR (END):14
DUPLICATE PREFERENCE IS 'EMBASE, CAPLUS'
KEEP DUPLICATES FROM MORE THAN ONE FILE? Y/(N):n
PROCESSING COMPLETED FOR L4
L5 8 DUPLICATE REMOVE L4 (1 DUPLICATE REMOVED)

=> d l5 1-8 bib ab

L5 ANSWER 1 OF 8 CAPLUS COPYRIGHT 2006 ACS on STN
AN 2006:184858 CAPLUS
TI Three-dimensional structure of glucodextranase, a glycoside
hydrolase
family 15 enzyme
AU Mizuno, Masahiro; Tonozuka, Takashi; Ichikawa, Kazuhiro;
Kamitori,
Shigehiro; Nishikawa, Atsushi; Sakano, Yoshiyuki
CS Department of Applied Science, Tokyo University of Agriculture
and
Technology, 3-5-8, Saiwai-cho, Fuchu, Tokyo, 183-8509, Japan
SO Biologia (Bratislava, Slovakia) (2005), 60(Suppl. 16), 171-176
CODEN: BLOAAO; ISSN: 0006-3088
PB Slovak Academy of Sciences
DT Journal
LA English
AB A glucodextranase (GDase) from Arthrobacter globiformis I42
hydrolyzes
to α -1,6-glucosidic linkages of dextran from the non-reducing end
to produce β -D-glucose. Here, we **review** the crystal
structures of GDase of the unliganded form and the complex with
acarbose.

The structure of GDase is composed of four domains, N, A, B, and C.

Domain A forms an (α/α)₆-barrel structure and domain N consists of 17 antiparallel β -strands. Both domains are conserved in

bacterial glucoamylases (GAs) and appear to be mainly concerned with

catalytic activity. The structure of GDase complexed with acarbose

revealed that the positions and orientations of the residues at subsites

-1 and +1 are nearly identical for GDase and GA; however, the residues

corresponding to subsite +2, which form the entrance of the substrate-binding pocket, and the position of the open space and constriction of GDase are different from those of GAs. On the other hand,

domains B and C are not found in bacterial GAs. The primary structure of

domain C is homologous with the surface layer homol. domain of pullulanases, and the three-dimensional **structure** of domain C resembles the **carbohydrate-binding domain** of some glycohydrolases. The hydrophobicity of domain B is higher than that

of the other three domains. These findings suggest that domains B and C

serve as cell wall anchors and contribute to the effective degradation of

dextran at the cell surface.

RE.CNT 35 THERE ARE 35 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 2 OF 8 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2005:503905 CAPLUS

DN 143:92803

TI X-ray crystallographic study of glucodextranase from a gram-positive

bacterium, *Arthrobacter globiformis* I42

AU Mizuno, Masahiro; Tonozuka, Takashi; Ichikawa, Kazuhiro; Kamitori,

Shigehiro; Nishikawa, Atsushi; Sakano, Yoshiyuki

CS United. Grad. Sch. Agric. Sci., Tokyo Univ. Agric. Technol., Fuchu,

183-8509, Japan

SO Journal of Applied Glycoscience (2005), 52(2), 145-151
CODEN: JAGLFX; ISSN: 1344-7882

PB Japanese Society of Applied Glycoscience

DT Journal; General Review

LA English

AB A **review**. Glucodextranase (GDase) hydrolyzes α -1,6-glucosidic linkages of dextran from the non-reducing end to

produce β -D-glucose. GDase is classified under GH15, whose major member is glucoamylase (GA) that hydrolyzes α -1,4-glucosidic linkages of starch. We have cloned a GDase gene from the gram-pos. bacterium *Arthrobacter globiformis* I42 and determined the crystal structure at 2.42-Å resolution. The structure of GDase is composed of four domains N, A, B and C. Domain N consists of 17 antiparallel β -strands and domain A forms an $(\alpha/\alpha)_6$ barrel structure, which is conserved between GAs. Furthermore, the complex structure with acarbose was also determined at 2.42-Å resolution. The structure of GDase complexed with acarbose revealed that the positions and orientations of the residues at subsites -1 and +1 are nearly identical for GDase and GA; however, Glu380 and Trp582 located at subsite +2, which form the entrance of the catalytic pocket, and the position of the open space and constriction of GDase are different from those of GAs. On the other hand, domains B and C are not found in GAs. The primary structure of domain C is homologous with the surface layer homol. (SLH) of pullulanases from Gram-pos. bacteria, and the three-dimensional **structure** of domain C resembles the **carbohydrate-binding domain** of some glycohydrolases. The hydrophobicity of domain B is higher than that of the other three domains. These findings suggest that domains B and C serve as cell wall anchors and contribute to the effective degradation of dextran at the cell surface.

L5 ANSWER 3 OF 8 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2003:830926 CAPLUS

DN 140:58465

TI Cellulose-binding domains: Tools for innovation in cellulosic fiber

production and modification

AU Quentin, Michael; van der Valk, Henry; van Dam, Jan; de Jong, Ed

CS Department of Fibre and Paper Technology, ATO BV, Wageningen, 6700 AA,

Neth.

SO ACS Symposium Series (2003), 855(Applications of Enzymes to Lignocellulosics), 132-155

CODEN: ACSMC8; ISSN: 0097-6156

PB American Chemical Society
DT Journal; General Review
LA English
AB A **review**. Plant cell walls are composed of cellulose, nature's most abundant macromol., and therefore represent a renewable resource of special tech. importance. Cellulose degrading enzymes involved in plant cell wall loosening (expansins), or produced by plant pathogenic microorganisms (cellulases), share similarities favoring the degradation of this highly crystalline substrate. Most of the cellulases and cell wall loosening expansins share a multi-domain **structure**, which includes a cellulose-binding domain (**CBD**). CBDs possess the intrinsic ability to strongly and specifically bind to cellulose. CBDs may be applied to engineer hybrid enzymes able to bind to cellulose on one end, and to display enzymic or chemical reactivity on the other, providing innovative solns. to modify cellulosic surfaces or to immobilize biocatalysts on it. In transgenic plants, CBDs influence polysaccharide synthesis and their assembly in the cell wall. Therefore, CBDs represent biotechnol. tools to modify cellulosic fibers either during their growth or during post harvest processing.

RE.CNT 125 THERE ARE 125 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 4 OF 8 CAPLUS COPYRIGHT 2006 ACS on STN
AN 2000:779465 CAPLUS
DN 134:38967
TI Phage display of cellulose binding domains for biotechnological application
AU Benhar, Itai; Tamarkin, Aviva; Marash, Lea; Berdichevsky, Yevgeny; Yaron, Sima; Shoham, Yuval; Lamed, Raphael; Bayer, Edward A.
CS Department of Molecular Microbiology and Biotechnology, The George S. Wise Faculty of Life Sciences, Tel-Aviv University, Ramat Aviv, Israel
SO ACS Symposium Series (2000), 769(Glycosyl Hydrolases for Biomass Conversion), 168-189
CODEN: ACSMC8; ISSN: 0097-6156
PB American Chemical Society
DT Journal; General Review
LA English
AB A **review** with 65 refs. In recent years, cellulose-binding domains (CBDs) derived from the cellulolytic systems of cellulose-degrading microorganisms have become a focal point of attention

for a wide range of biotechnol. applications. The low cost and availability of cellulose matrixes have rendered CBDs attractive as

affinity tags for the purification and immobilization of a plethora of proteins. Intensive studies of cellulose degradation pathways and the

identification of components of the cellulose-degrading machinery have

contributed significantly to our understanding of the **structure** and function of **CBDs**. The time is now ripe to utilize engineered CBDs to improve existing applications and to devise novel ones.

Here we describe our recent results of expts. where the *Clostridium*

thermocellum scaffoldin CBD was genetically engineered for such purposes.

We describe the development of a novel phage display system where the C.

thermocellum CBD is displayed as a fusion protein with single-chain

antibodies. Our system is a filamentous (M13) phage display system that

enables the efficient isolation and.

RE.CNT 65 THERE ARE 65 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 5 OF 8 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2000:710413 CAPLUS

DN 134:38648

TI The structure and function of cellulose-binding domain of cellulase

AU Wang, Tianhong; Wang, Chunhui; Gao, Peiji

CS The State Key Laboratory of Microbial Technology, Shandong University,

Jinan, 250100, Peop. Rep. China

SO Shengwu Gongcheng Jinzhan (2000), 20(2), 37-40

CODEN: SGJHA2; ISSN: 1003-3505

PB Zhongguo Kexueyuan Wenxian Qingbao Zhongxin

DT Journal; General Review

LA Chinese

AB A discussion and **review** with 20 refs. Most cellulases consist of catalytic domains and cellulose binding domains(CBDs), which can bind

to cellulose and are conserved in some amino acid sequences.

Cellulose

binding domains improve the binding and facilitate the activity of

catalytic domains on the insol. substrate, but not on soluble substrate. The

results of investigations on structure and function, and subsequent

mutagenesis of the CBDs indicated that CBDs rely on several aromatic amino acids for binding to the cellulose surfaces. Some experiment results showed that CBDs of exoglucanases are able to disrupt the crystalline cellulose, facilitate the enzymic degradation of cellulose. The structural domain as CBDs has been successfully used in purification and immobilization of numerous examples of fusion proteins. The improved understanding of the **structure** and function of **CBDs** are significant to understanding of enzymic functionary mechanism, and the development of cellulase biotechnol.

L5 ANSWER 6 OF 8 CAPLUS COPYRIGHT 2006 ACS on STN

AN 1998:330997 CAPLUS

DN 129:105755

TI The structure and function of cellulose binding domains

AU Boraston, A.; Bray, M.; Brun, E.; Creagh, A. L.; Gilkes, N. R.; Guarna, M.

M.; Jervis, E.; Johnson, P.; Kormos, J.; McIntosh, L.; Mclean, B. W.;

Sandercock, L. E.; Tomme, P.; Haynes, C. A.; Warren, R. A. J.; Kilburn, D.

G.

CS Protein Engineering Network of Centres of Excellence, Biotechnology

Laboratory, University of British Columbia, Vancouver, V6T 1Z3, Can.

SO Special Publication - Royal Society of Chemistry (1998), 219(Carbohydrates

from Trichoderma Reesei and Other Microorganisms), 139-146

CODEN: SROCD0; ISSN: 0260-6291

PB Royal Society of Chemistry

DT Journal; General Review

LA English

AB A **review** with 15 refs. More than 180 putative cellulose-binding domains (CBDs) have been identified and grouped into 13 families based on

amino acid sequences. They vary in length from 33 to 240 amino acids.

The **structures** of **CBDs** from five different families are now available. They all are anti-parallel β strand polypeptides.

The authors have studied the properties of Family II and Family IV CBDs of

β -glycanases from the cellulolytic bacterium Cellulomonas fimi in

detail.

RE.CNT 15 THERE ARE 15 CITED REFERENCES AVAILABLE FOR THIS RECORD

ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 7 OF 8 EMBASE COPYRIGHT (c) 2006 Elsevier B.V. All rights reserved on STN DUPLICATE 1

AN 94168662 EMBASE

DN 1994168662

TI The Clostridium cellulovorans cellulosome.

AU Doi R.H.; Goldstein M.; Hashida S.; Park J.-S.; Takagi M.

CS Molecular/Cellular Biology Section, University of California, Davis, CA 95616, United States

SO Critical Reviews in Microbiology, (1994) Vol. 20, No. 2, pp. 87-93. .

ISSN: 1040-841X CODEN: CRVMAC

CY United States

DT Journal; General Review

FS 004 Microbiology

LA English

SL English

ED Entered STN: 29 Jun 1994

Last Updated on STN: 29 Jun 1994

AB The Clostridium cellulovorans cellulosome is comprised of a large, nonenzymatic scaffolding protein called the cellulose binding protein A (CbpA) and a number of endoglucanases/xylanases. The CbpA contains several functional domains, including a signal peptide, a cellulose binding domain (CBD), a hydrophilic domain (HLD) present four times, and a hydrophobic domain (HBD) present nine times. The functions of the domains were studied by the construction of minigenes containing the putative functional domains and by expression of the minigenes in Escherichia coli. The purified product of the CBD was able to bind to various crystalline forms of cellulose and chitin with a K(d) of 1 μ M. The binding capacity for CBD was a function of the crystallinity of the cellulose sample. Furthermore, the binding of CBD to Avicel was not inhibited by cellobiose or carboxymethylcellulose, suggesting that the CBD binding target was a three-dimensional **structure** found only in crystalline forms of cellulose. The HBD was tested for its ability to bind endoglucanases by an interaction Western as well as a sandwich enzyme immunoassay technique. The HBD was able to bind both EngB and EngD,

indicating that the HBD contained an endoglucanase binding domain (EBD).

Because there are nine EBD domains, it is possible that CbpA can bind up

to nine endoglucanases. The role of the HLDs remains elusive. The data

indicate that the cellulosome is a complex enzyme containing a scaffolding

protein (CbpA) to which is attached a number of endoglucanase molecules.

This arrangement allows the complex to bind and degrade crystalline

cellulose, which resists degradation by the free forms of cellulosomal

endoglucanases.

L5 ANSWER 8 OF 8 CAPLUS COPYRIGHT 2006 ACS on STN

AN 1994:263906 CAPLUS

DN 120:263906

TI Lectins. Structures and lectin-sugar interactions

AU Yamamoto, Kazuo

CS Fac. Pharm. Sci., Univ. Tokyo, Tokyo, 113, Japan

SO Maku (1994), 19(1), 40-7

CODEN: MAKUD9; ISSN: 0385-1036

DT Journal; General Review

LA Japanese

AB A **review**, with 42 refs. Topics include: classification of lectins, variety of sugar chain **structures**, determination of **carbohydrate-binding domains** of legume

lectins, chemical synthesis of a lactose-binding domain (nonapeptide),

construction of a chimeric lectin replaced with another nonapeptide from a

lectin having a different carbohydrate-binding specificity, and conformation of the carbohydrate-binding domain. The anal. of binding

specificity of the chimeric lectin shows the presence of a variable-binding region in legume lectins that determine their carbohydrate-binding specificity. The result suggests the presence of a

variable-binding region in legume lectins that determine their carbohydrate-binding specificity.

=> s (glycosyl hydrolase 74) or (gh74)

L6 10 (GLYCOSYL HYDROLASE 74) OR (GH74)

=> s 16 (6A) structure

L7 0 L6 (6A) STRUCTURE